

GRADED POSETS ZETA MATRIX FORMULA

Andrzej Krzysztof Kwaśniewski

Member of the Institute of Combinatorics and its Applications
 High School of Mathematics and Applied Informatics
 Kamienna 17, PL-15-021 Białystok, Poland
 e-mail: kwandr@gmail.com

Abstract: The way to arrive at formula of zeta matrix for any graded posets with the finite set of minimal elements is delivered following [1]. This is being achieved via adjacency and zeta matrix description of bipartite digraphs chains - the representatives of graded posets. The bipartite digraphs elements of such chains amalgamate to form corresponding cover relation graded poset digraphs with corresponding adjacency matrices being amalgamated throughout natural join as special adequate database operation. The colligation of reachability and connectivity with the presented description is made explicit. The special posets encoded via KoDAGs directed acyclic graphs as cobeb posets Hasse diagrams are recognized as an example of differential posets sub-family. As on the 01.01.2009 one reminisce 261-th anniversary of death of Johann Bernoulli the First this Sylvester Night article is to commemorate this date.

Key Words: graded digraphs, differential posets, natural join, umbral calculus

AMS Classification Numbers: 06A06 ,05B20, 05C75 , 05A30

affiliated to The Internet Gian-Carlo Rota Polish Seminar:

http://ii.uwb.edu.pl/akk/sem/sem_rota.htm

1 Preliminaries: notation and terminology.

1.1. We shall try to keep track of NIST Dictionary of Algorithms and Data Structures terminology. Abbreviation: directed acyclic graph = DAG. Note: The transitive closure of a directed acyclic graph or DAG is the reachability relation of the DAG and a strict partial order.

1.2. The following convention scheme is adopted: directed graph representatives of binary relations scheme are:

$$\begin{aligned}
 &\text{bipartite digraph representative } D_R = (A \times A, R) \leftrightarrow R \subseteq A \times A \\
 &\equiv \text{"just" digraph representative } D(R) \equiv D_R = (A, E), E \leftrightarrow R \subseteq A \times A, \\
 &\text{bipartite digraph representative } D_R = (A \times B, R) \leftrightarrow R \subseteq A \times B
 \end{aligned}$$

1.3 A directed path, is an oriented simple path with all arcs of the same direction i.e. all internal nodes have in- and out-degrees equal one.

Comment 1.

"A directed path is a natural join of arcs that thus form a chain of vertices",
 "A chain of coded data objects is a natural join of their subsequent pairs".

Anticipated: coded data objects = relations (with varying arity allowed), binary relations, bipartite digraphs, adjacency matrices (of graphs or digraphs): [1], [2].

Comment 2.

Because of immense number of applications of digraphs beyond mathematics - frequently successfully done by non-mathematicians - it happens sometimes that various names are given for the same notions and different names for the same objects. Let us recall and/or establish some of them.

Recall

Reachability is the ability reach some other vertex from a given vertex in a directed graph. For a directed graph $D = (\Phi, E)$, $E \subseteq \Phi \times \Phi$ the reachability relation of D is its transitive closure of E , i.e. the set of all ordered pairs (s, t) of vertices in Φ for which there exist vertices $v_0 = s, v_1, \dots, v_e = t$ such that $(v_{i-1}, v_i) \in E$ for all $1 \leq i \leq e$.

We define here (for directed graphs - more than nontrivial) the reachability = connectivity partial order relation R over the nodes of the DAG as such that xRy iff there exists a directed path from x to y .

1.4. (relations set sum)

$$\begin{aligned} \text{reachability} \cup \text{reflexibility} &= \text{reflexive reachability,} \\ \text{reachability} &= \text{connectivity,} \end{aligned}$$

This means that:

1.4.1 The reachability = connectivity relation is $\mathbf{R}^\infty = \bigcup_{k>0} R^k = \text{transitive closure of } \mathbf{R}$, i.e.

$$\mathbf{R}^\infty = R^1 \cup R^2 \cup \dots \cup R^n \cup \dots \Leftrightarrow A(R^\infty) = A(R)^{\odot 1} \vee A(R)^{\odot 2} \vee \dots \vee A(R)^{\odot n} \vee \dots,$$

where $A(\mathbf{R})$ is the Boolean adjacency matrix of the relation $\mathbf{R}$ simple digraph and $\odot$ stays for Boolean product.

The **reflexive** reachability relation $\zeta(\mathbf{R}) \equiv \mathbf{R}^*$ is defined as

$$\begin{aligned} \mathbf{R}^* &= R^0 \cup R^1 \cup R^2 \cup \dots \cup R^n \cup \dots \bigcup_{k \geq 0} R^k = \mathbf{R}^\infty \cup \mathbf{I}_A = \\ &= \text{transitive and reflexive closure of } \mathbf{R} \Leftrightarrow \end{aligned}$$

$$\Leftrightarrow A(R^\infty) = A(R)^{\odot 0} \vee A(R)^{\odot 1} \vee A(R)^{\odot 2} \vee \dots \vee A(R)^{\odot n} \vee \dots$$

Comment 3. Colligate and identify $\zeta(\mathbf{R}) \equiv \mathbf{R}^*$ with incidence algebra zeta function and with zeta matrix of the poset associated to its Hasse digraph.

1.4.2. (Notation continued)

In what follows we shall use mathematical terms: reachability and reflexive reachability according to: put $R = \prec \cdot$ which is Hasse diagram i.e. cover relation digraph - notation. Then note the **schemes** below.

The partial order $\leq$ for locally finite poset $\Pi = \langle \Phi, \leq \rangle$ with respect to Π 's cover relation $\prec \cdot$ is:

1. **$<$ the connectivity relation in Hasse digraph i.e. digraph $D = \langle \Phi, \prec \cdot \rangle$.**

2. $\leq$ is the reflexive reachability relation in Hasse digraph i.e. digraph $D = \langle \Phi, \prec \cdot \rangle$.

Schemes:

$$\begin{aligned} \leq &= \prec \cdot^\infty = \text{connectivity of } \prec \cdot \\ \leq &= \prec \cdot^* = \text{reflexive reachability of } \prec \cdot \\ \prec \cdot^* &= \zeta(\prec \cdot). \end{aligned}$$

1.4.3. $\zeta(\prec \cdot)$ is also used to denote zeta matrix of the graded poset $P(D) = \Pi = \langle \Phi, \leq \rangle$ associated to $D = \langle \Phi, \prec \cdot \rangle$ [1,2] which is equivalent to say that $P(D) = \langle \Phi, \leq \rangle =$ transitive, reflexive closure of $D = \langle \Phi, \prec \cdot \rangle$.

1.5.

In order to get **complete** graded digraph connect any two vertices lying on consecutive levels with an arc keeping one direction - say - upwards (see KoDAG in [1,2]).

2 Natural join of adjacency matrices

2.1. Locally finite poset Π if fixed, is denominated by all its covering pairs and vice versa of course i.e.

$$\Pi = \langle \Phi, \leq \rangle \Leftrightarrow \langle \Phi, \prec \cdot \rangle,$$

as all properties of order origin follow from those of transitivity requirement - and vice versa of course. Specifically recall-note the equivalence of descriptions:

The complete graded poset $\Leftrightarrow$ The complete graded digraph.

The arcs of any digraph $G = \langle \Phi, E \rangle$ with no multiple edges stay automatically for arcs of cover relation $\prec \cdot$ in the corresponding poset $\Pi = \langle \Phi, \leq \rangle$ for which the digraph $G = \langle \Phi, E \rangle$ becomes Hasse diagram i.e. we have:

$$\begin{aligned} \leq &= \prec \cdot^* = \text{reflexive reachability of } \prec \cdot \\ \prec \cdot^* &\equiv (I - \prec \cdot)^{-1} \equiv \zeta(\prec \cdot) \end{aligned}$$

Unless differently stated, we shall identify a digraph with its adjacency matrix. In our context these are to be Hasse digraphs $D = \langle \Phi, \prec \cdot \rangle$ of graded posets $\Pi = \langle \Phi, \leq \rangle$.

There are three standard widely used prevalent encodings, three ways of portraying partially ordered sets $P(D) = \langle \Phi, \leq \rangle$: Hasse diagrams $D = \langle \Phi, \prec \cdot \rangle$; zeta matrices $\zeta(\leq)$; and cover matrices $\zeta(\prec \cdot)$. These matrices are of course the adjacency matrices of the corresponding digraphs $\langle \Phi, \leq \rangle$ and $\langle \Phi, \prec \cdot \rangle$. In the incidence algebra description of locally finite posets $\zeta(\leq)$ may be identified with the incidence function i.e. the characteristic function of the partial order $\leq$, (see [1] and references therein for the source cobweb posets examples of these objects). Here down we shall use adjacency and biadjacency nomenclature [1,2].

Examples of $\zeta(\leq)$

Let F denotes arbitrary natural numbers valued sequence. Let A_N be the Hasse matrix .i.e. adjacency matrix of cover relation $\prec \cdot$ digraph denominated by sequence N [1]. Then the zeta matrix $\zeta = (1 - \mathbf{A}_N)^{-1} \textcircled{\text{C}}$ for the denominated by $F = N$ cobweb poset is of the form [1]

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 1 & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 1 & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \mathbf{0} & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \dots \end{bmatrix}$$

Figure ζ_N . The incidence matrix ζ for the natural numbers i.e. N - cobweb poset

Note that the matrix ζ representing uniquely its corresponding cobweb poset does exhibits a staircase structure of zeros above the diagonal (see above, see below) which is characteristic to and characteristics of Hasse diagrams of **all cobweb posets** while for graded posets it is characteristic too - this time all together with additional zeros right to the staircase - zeros generated by the bi-adjacency matrices $\langle B_k \rangle_{k \geq 0}$ chain as described in Observation 4. . Another wards: it is the natural join $\oplus \rightarrow$ chain

$$\oplus \rightarrow_{k=0}^n B_k, \quad n \in N \cup \{\infty\}$$

which is equivalent to ζ function **characteristics** of any fixed F -denominated graded poset $(P, \leq)$.

$$\begin{bmatrix}
1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 1 & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & 1 & 1 & 1 & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \mathbf{0} & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \mathbf{0} & \dots \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots
\end{bmatrix}$$

Figure ζ_F . The matrix ζ for the Fibonacci cobweb poset associated to F -KoDAG Hasse digraph

Comment 4. The given F -denominated staircase zeros structure above the diagonal of zeta matrix ζ is the **unique characteristics** of its corresponding F -KoDAG Hasse digraphs.

2.2. The natural join condition.

The natural join operation is a binary operation like Θ operator in computer science denoted here by $\oplus \rightarrow$ symbol deliberately referring - in a quite reminiscent manner - to direct sum $\oplus$ of adjacency Boolean matrices and - as matter of fact and in effect - to direct the sum $\oplus$ of corresponding biadjacency [reduced] matrices of digraphs under natural join.

$\oplus \rightarrow$ is a natural operator for sequences construction. $\oplus \rightarrow$ operates on multi-ary relations according to the scheme: $(n + k)_{ary} \oplus \rightarrow (k + m)_{ary} = (n + k + m)_{ary}$ For example: $(1 + 1)_{ary} \oplus \rightarrow (1 + 1)_{ary} = (1 + 1 + 1)_{ary}$, binary $\oplus \rightarrow$ binary = ternary.

Accordingly an action of $\oplus \rightarrow$ on these multi-ary relations' digraphs adjacency matrices is to be designed soon in what follows.

Domain-Codomain F -sequence condition $\text{dom}(R_{k+1}) = \text{ran}(R_k)$, $k = 0, 1, 2, \dots [1]$.

Consider any natural number valued sequence $F = \{F_n\}_{n \geq 0}$. Consider then any chain of binary relations defined on pairwise disjoint finite sets with cardinalities appointed by F -sequence elements values. For that to start we specify at first a relations' domain-co-domain F -sequence.

Domain-Codomain F -sequence $(|\Phi_n| = F_n)$

$$\Phi_0, \Phi_1, \dots, \Phi_i, \dots \quad \Phi_k \cap \Phi_n = \emptyset \text{ for } k \neq n, |\Phi_n| = F_n; \quad i, k, n = 0, 1, 2, \dots$$

Let $\Phi = \bigcup_{k=0}^n \Phi_k$ be the corresponding ordered partition [anticipating - Φ is the vertex set of $D = (\Phi, \prec \cdot)$ and its transitive, reflexive closure $(\Phi, \leq)$]. Impose $\text{dom}(R_{k+1}) = \text{ran}(R_k)$ condition , $k \in N \cup \{\infty\}$. What we get is binary relations chain.

Definition 1 (*Relation's chain*) Let $\Phi = \bigcup_{k=0}^n \Phi_k$, $\Phi_k \cap \Phi_n = \emptyset$ for $k \neq n$ be the ordered partition of the set Φ . Let a sequence of binary relations be given such that

$$R_0, R_1, \dots, R_i, \dots, R_{i+n}, \dots, \quad R_k \subseteq \Phi_k \times \Phi_{k+1}, \quad \text{dom}(R_{k+1}) = \text{ran}(R_k).$$

Then the sequence $\langle R_k \rangle_{k \geq 0}$ is called *natural join (binary) relation's chain*.

Extension to varying arity relations' natural join chains is straightforward.

As necessarily $\text{dom}(R_{k+1}) = \text{ran}(R_k)$ for relations' natural join chain any given binary relation's chain is not just a sequence therefore we use "link to link " notation for $k, i, n = 1, 2, 3, \dots$ ready for relational data basis applications:

$R_0 \oplus \rightarrow R_1 \oplus \rightarrow \dots \oplus \rightarrow R_i \oplus \rightarrow \dots \oplus \rightarrow R_{i+n}, \dots$ is an F -chain of binary relations

where $\oplus \rightarrow$ denotes natural join of relations as well as both natural join of their bipartite digraphs and the natural join of their representative adjacency matrices (see [1,2]).

Relation's F -chain naturally represented by [identified with] the chain of theirs **bipartite digraphs**

$$\begin{aligned} R_0 \oplus \rightarrow R_1 \oplus \rightarrow \dots \oplus \rightarrow R_i \oplus \rightarrow \dots \oplus \rightarrow R_{i+n}, \dots &\Leftrightarrow \\ \Leftrightarrow B_0 \oplus \rightarrow B_1 \oplus \rightarrow \dots \oplus \rightarrow B_i \oplus \rightarrow \dots \oplus \rightarrow B_{i+n}, \dots \end{aligned}$$

results in **F -partial ordered set** $\langle \Phi, \leq \rangle$ with its Hasse digraph representation looking like specific "cobweb" image (for cobweb posets portraits see [1] and references therein and see also cobwebs in action on <http://www.faces-of-nature.art.pl/cobwebposets.html>).

2.3 Partial order $\leq$.

The partial order relation $\leq$ in the set of all points-vertices is determined uniquely by the above equivalent F - chains. Let $x, y \in \Phi = \bigcup_{k=0}^n \Phi_k$ and let $k, i = 0, 1, 2, \dots$. Then

$$(1) \quad x \leq y \Leftrightarrow \forall_{x \in \Phi} : x \leq x \vee \Phi_i \ni x < y \in \Phi_{i+k} \text{ iff } x(R_i \odot \dots \odot R_{i+k-1})y$$

where " $\odot$ " stays for [Boolean] composition of binary relations.

Relation ($\leq$) is defined equivalently :

$x \leq y$ in $(\Phi, \leq)$ iff either $x = y$ or there exist a directed path from x to y ; $x, y \in \Phi$.

Let now $R_k = \Phi_k \times \Phi_{k+1}, k \in N \cup \{0\}$. For "historical" reasons [1] we shall call such partial ordered set $\Pi = \langle \Phi, \leq \rangle$ the **cobweb poset** as theirs Hasse digraph representation looks like specific "cobweb" image.

2.4. The natural join $\oplus \rightarrow$ operation and the natural join of matrices satisfying the natural join condition

We define here the adjacency matrices representation of the natural join $\oplus \rightarrow$ operation.

The adjacency matrix $\mathbf{A}$ of a bipartite graph with **biadjacency** = reduced adjacency matrix $\mathbf{B}$ is given by

$$\mathbf{A} = \begin{pmatrix} 0 & \mathbf{B} \\ \mathbf{B}^T & 0 \end{pmatrix}.$$

Definition 2 The adjacency matrix $\mathbf{A}[D]$ of a bipartite **digraph** $D(R) = (P \cup L, E \subseteq P \times L)$ with biadjacency matrix $\mathbf{B}$ is given by [1]

$$\mathbf{A}[D] = \begin{pmatrix} 0_{k,k} & \mathbf{B}(k \times m) \\ 0_{m,k} & 0_{m,m} \end{pmatrix}.$$

where $k = |P|$, $m = |L|$.

Note: biadjacency and cover relation $\prec \cdot$ matrix for bipartite digraphs coincide. By extension - we shall call cover relation $\prec \cdot$ matrix also the biadjacency matrix.

Convention 1 $S \odot R$ = composition of binary relations S and $R \Leftrightarrow \mathbf{B}_R \odot \mathbf{B}_S = \mathbf{B}_R \odot \mathbf{B}_S$ where $(|V| = k, |W| = m) \mathbf{B}_R(k \times m) \equiv \mathbf{B}_R$.

$\mathbf{B}_R$ is the $(k \times m)$ **biadjacency** [or another name: **reduced** adjacency] matrix of the bipartite relations' R digraph $B(R)$ and $\odot$ apart from relations composition denotes also Boolean multiplication of these rectangular biadjacency Boolean matrices $\mathbf{B}_R, \mathbf{B}_S$. What is their form? The answer is in the block structure of the standard square $(n \times n)$ adjacency matrix $\mathbf{A}[D(R)]; n = k + m$. The form of standard square adjacency matrix $\mathbf{A}[G(R)]$ of bipartite digraph $D(R)$ has the following apparently recognizable block reduced structure: $[O_{s \times s}$ stays for $(k \times m)$ zero matrix]

$$\mathbf{A}[D(R)] = \begin{bmatrix} O_{k \times k} & \mathbf{A}_R(k \times m) \\ O_{m \times k} & O_{m \times m} \end{bmatrix}$$

Let $D(S) = (W(S) \cup T(S), E(S)); W \cap T = \emptyset, E(S) \subseteq W \times T; (|W| = m, |T| = s)$; hence

$$\mathbf{A}[D(S)] = \begin{bmatrix} O_{m \times m} & \mathbf{A}_S(m \times s) \\ O_{s \times m} & O_{s \times s} \end{bmatrix}$$

Definition 3 (natural join condition) The ordered pair of matrices $\langle \mathbf{A}_1, \mathbf{A}_2 \rangle$ is said to satisfy the natural join condition iff they have the block structure of $\mathbf{A}[D(R)]$ and $\mathbf{A}[D(S)]$ as above i.e. iff they might be identified accordingly : $\mathbf{A}_1 = \mathbf{A}[D(R)]$ and $\mathbf{A}_2 = \mathbf{A}[D(S)]$.

Correspondingly if two given digraphs G_1 and G_2 are such that their adjacency matrices $\mathbf{A}_1 = \mathbf{A}[G_1]$ and $\mathbf{A}_2 = \mathbf{A}[G_2]$ do satisfy the natural join condition we shall say that G_1 and G_2 satisfy the natural join condition. For matrices satisfying the natural join condition one may define what follows.

First we define the **Boolean reduced** or **natural join composition** $\odot \rightarrow$ and secondly the natural join $\oplus \rightarrow$ of adjacent matrices satisfying the natural join condition.

Definition 4 ($\odot \rightarrow$ composition)

$$\mathbf{A}[D(R \odot S)] =: \mathbf{A}[D(R)] \odot \rightarrow \mathbf{A}[D(S)] = \begin{bmatrix} O_{k \times k} & \mathbf{A}_R \odot_S (k \times s) \\ O_{s \times k} & O_{s \times s} \end{bmatrix}$$

where $\mathbf{A}_R \odot_S (k \times s) = \mathbf{A}_R(k \times m) \odot \mathbf{A}_S(m \times s)$.

according to the scheme:

$$[(k + m) \times (k + m)] \odot \rightarrow [(m + s) \times (m + s)] = [(k + s) \times (k + s)].$$

Comment 5. The adequate projection makes out the intermediate, joint in common $\text{dom}(S) = \text{rang}(R) = W$, $|W| = m$.

The above Boolean reduced composition $\odot \rightarrow$ of adjacent matrices technically reduces then to the calculation of just Boolean product of the **reduced** rectangular adjacency matrices of the bipartite relations' graphs.

We are however now in need of the Boolean natural join product $\oplus \rightarrow$ of adjacent matrices already announced at the beginning of this presentation. Let us now define it.

As for the **natural join** notion we aim at the morphism correspondence:

$$S \oplus \rightarrow R \Leftrightarrow M_{S \oplus \rightarrow R} = M_R \oplus \rightarrow M_S$$

where $S \oplus \rightarrow R$ = natural join of binary relations S and R while $M_{S \oplus \rightarrow R} = M_R \oplus \rightarrow M_S$ = natural join of standard square adjacency matrices (with customary convention: $M[G(R)] \equiv M_R$ adapted). Attention: recall here that the natural join of the above binary relations $R \oplus \rightarrow S$ is the ternary relation - and thus one results in k -ary relations if with more factors undergo the $\oplus \rightarrow$ product. As a matter of fact $\oplus \rightarrow$ **operates on multi-ary relations according to the scheme:**

$$(n + k)_{ary} \oplus \rightarrow (k + m)_{ary} = (n + k + m)_{ary}.$$

For example: $(1 + 1)_{ary} \oplus \rightarrow (1 + 1)_{ary} = (1 + 1 + 1)_{ary}$, $binary \oplus \rightarrow binary = ternary$.

Technically - the natural join of the k -ary and n -ary relations is defined accordingly the same way via $\oplus \rightarrow$ natural join product of adjacency matrices - the adjacency matrices of these relations' Hasse digraphs.

With the notation established above we finally define the natural join $\oplus \rightarrow$ of two adjacency matrices as follows:

Definition 5 (natural join $\oplus \rightarrow$ of matrices) .

$$\begin{aligned}
A[D(R \oplus \rightarrow S)] &=: A[D(R)] \oplus \rightarrow A[D(S)] = \\
&= \begin{bmatrix} O_{k \times k} & A_R(k \times m) \\ O_{m \times k} & O_{m \times m} \end{bmatrix} \oplus \rightarrow \begin{bmatrix} O_{m \times m} & A_S(m \times s) \\ O_{s \times m} & O_{s \times s} \end{bmatrix} = \\
&= \begin{bmatrix} O_{k \times k} & A_R(k \times m) & O_{k \times s} \\ O_{m \times k} & O_{m \times m} & A_S(m \times s) \\ O_{s \times k} & O_{s \times m} & O_{s \times s} \end{bmatrix}
\end{aligned}$$

Comment 6. The adequate projection used in natural join operation leaves one copy of the joint in common "intermediate" submatrix $O_{m \times m}$ and consequently leaves one copy of "intermediate" joint in common m according to the scheme:

$$[(k + m) \times (k + m)] \oplus \rightarrow [(m + s) \times (m + s)] = [(k + m + s) \times (k + m + s)].$$

2.5. The biadjacency i.e cover relation $\prec \cdot$ matrices of the natural join of adjacency matrices.

Denote with $B(A)$ the biadjacency i.e cover relation $\prec \cdot$ matrix of the adjacency matrix A .

Note: biadjacency and cover relation $\prec \cdot$ matrix for bipartite digraphs coincide. By extension - we shall call cover relation $\prec \cdot$ matrix the biadjacency matrix too, as for any graded digraph with more than one level we might represent it as a partition into two independent sets, though it is more natural to see it as the natural join of the sequence of bipartite digraphs

Let $A(G)$ denotes the adjacency matrix of the digraph G , for example a dibiclique relation digraph. Let $A(G_k)$, $k = 0, 1, 2, \dots$ be the sequence adjacency matrices of the sequence G_k , $k = 0, 1, 2, \dots$ of digraphs. Let us identify $B(A) \equiv B(G)$ as a convention.

Definition 6 (digraphs natural join) Let digraphs G_1 and G_2 satisfy the natural join condition. Let us make then the identification $A(G_1 \oplus \rightarrow G_2) \equiv A_1 \oplus \rightarrow A_2$ as definition. The digraph $G_1 \oplus \rightarrow G_2$ is called the digraphs natural join of digraphs G_1 and G_2 . Note that the order is essential.

We observe at once what follows.

Observation 1

$$B(G_1 \oplus \rightarrow G_2) \equiv B(A_1 \oplus \rightarrow A_2) = B(A_1) \oplus B(A_2) \equiv B(G_1) \oplus B(G_2)$$

Comment 7. The Observation 1 justifies the notation $\oplus \rightarrow$ for the natural join of relations digraphs and equivalently for the natural join of their adjacency matrices and equivalently for the natural join of relations that these are faithful representatives of. Recall: $B(A)$ is the biadjacency i.e cover relation $\prec \cdot$ matrix of the adjacency matrix A .

Note: biadjacency and cover relation $\prec \cdot$ matrix for bipartite digraphs coincide. Recall that by extension - we call cover relation $\prec \cdot$ matrix the biadjacency matrix too.

As a consequence we have.

Observation 2

$$B(\oplus_{i=1}^n G_i) \equiv B[\oplus_{i=1}^n A(G_i)] = \oplus_{i=1}^n B[A(G_i)] \equiv \text{diag}(B_1, B_2, \dots, B_n) =$$

$$= \begin{bmatrix} B_1 & & & \\ & B_2 & & \\ & & B_3 & \\ & \dots & \dots & \dots \\ & & & B_n \end{bmatrix},$$

or equivalently

$$\kappa \equiv \chi(\prec \cdot) = \chi(\oplus_{i=1}^n \prec \cdot_i) \equiv$$

$$\equiv \begin{bmatrix} 0 & B_1 & & & \\ & 0 & B_2 & & \\ & & 0 & B_3 & \\ & \dots & \dots & \dots & \\ & & & 0 & B_n \\ & & & & 0 \end{bmatrix},$$

$$n \in N \cup \{\infty\}$$

2.6. The formula of zeta matrix for graded posets with the finite set of minimal elements i.e for F -graded posets

Any graded poset with the finite set of minimal elements is an F - sequence denominated sub-poset of its corresponding cobweb poset. The Observation 2 supplies the simple recipe for the biadjacency (reduced adjacency) matrix of Hasse digraph coding any given graded poset with the finite set of minimal elements. The recipe for zeta matrix is then standard. We illustrate this by the source example; the source example as the adjacency matrices i.e zeta matrices of any given graded poset with the finite set of minimal elements are sub-matrices of their corresponding cobweb posets and as such have the same block matrix structure.

The explicit expression for zeta matrix ζ_F of **cobweb posets** via known blocks of zeros and ones for arbitrary natural numbers valued F - sequence was given in [1] due to more than mnemonic efficiency of the up-side-down notation being applied (see [1] and references therein). With this notation inspired by Gauss and replacing k - natural numbers with " k_F " numbers one gets

$$\mathbf{A}_F = \begin{bmatrix} 0_{1_F \times 1_F} & I(1_F \times 2_F) & 0_{1_F \times \infty} & & \\ 0_{2_F \times 1_F} & 0_{2_F \times 2_F} & I(2_F \times 3_F) & 0_{2_F \times \infty} & \\ 0_{3_F \times 1_F} & 0_{3_F \times 2_F} & 0_{3_F \times 3_F} & I(3_F \times 4_F) & 0_{3_F \times \infty} \\ 0_{4_F \times 1_F} & 0_{4_F \times 2_F} & 0_{4_F \times 3_F} & 0_{4_F \times 4_F} & I(4_F \times 5_F) & 0_{4_F \times \infty} \\ \dots & etc & \dots & and so on & \dots & \end{bmatrix}$$

and

$$\zeta_F = \exp_{\odot}[\mathbf{A}_F] \equiv (1 - \mathbf{A}_F)^{-1 \odot} \equiv I_{\infty \times \infty} + \mathbf{A}_F + \mathbf{A}_F^{\odot 2} + \dots =$$

$$= \begin{bmatrix} I_{1_F \times 1_F} & I(1_F \times \infty) & & & \\ O_{2_F \times 1_F} & I_{2_F \times 2_F} & I(2_F \times \infty) & & \\ O_{3_F \times 1_F} & O_{3_F \times 2_F} & I_{3_F \times 3_F} & I(3_F \times \infty) & \\ O_{4_F \times 1_F} & O_{4_F \times 2_F} & O_{4_F \times 3_F} & I_{4_F \times 4_F} & I(4_F \times \infty) \\ \dots & etc & \dots & and\ so\ on & \dots \end{bmatrix}$$

where $I(s \times k)$ stays for $(s \times k)$ matrix of ones i.e. $[I(s \times k)]_{ij} = 1; 1 \leq i \leq s, 1 \leq j \leq k$. and $n \in N \cup \{\infty\}$

Observation 3 Let us denote by $\langle \Phi_k \rightarrow \Phi_{k+1} \rangle$ (see the authors papers quoted) the di-bicliques denominated by subsequent levels Φ_k, Φ_{k+1} of the graded F -poset $P(D) = (\Phi, \leq)$ i.e. levels Φ_k, Φ_{k+1} of its cover relation graded digraph $D = (\Phi, \prec \cdot)$ [Hasse diagram]. Then

$$B(\oplus_{k=1}^n \langle \Phi_k \rightarrow \Phi_{k+1} \rangle) = \text{diag}(I_1, I_2, \dots, I_n) =$$

$$= \begin{bmatrix} I(1_F \times 2_F) & & & & \\ & I(2_F \times 3_F) & & & \\ & & I(3_F \times 4_F) & & \\ & & \dots & & \\ & & & I(n_F \times (n+1)_F) & \end{bmatrix}$$

where $I_k \equiv I(k_F \times (k+1)_F)$, $k = 1, \dots, n$ and where - recall - $I(s \times k)$ stays for $(s \times k)$ matrix of ones i.e. $[I(s \times k)]_{ij} = 1; 1 \leq i \leq s, 1 \leq j \leq k$. and $n \in N \cup \{\infty\}$.

The recipe for any F -denominated i.e. the recipe for **any graded poset** with a finite minimal elements set is supplied via the following observation.

Observation 4 Consider bigraphs' chain **obtained from** the above di-bicliques' chain via deleting or no arcs making thus [if deleting arcs] some or all of the di-bicliques $\langle \Phi_k \rightarrow \Phi_{k+1} \rangle$ not di-bicliques; denote them as G_k . Let $B_k = B(G_k)$ denotes their biadjacency matrices correspondingly. Then for any such F -denominated chain [hence any chain] of bipartite digraphs G_k the general formula is:

$$B(\oplus_{i=1}^n G_i) \equiv B[\oplus_{i=1}^n A(G_i)] = \oplus_{i=1}^n B[A(G_i)] \equiv \text{diag}(B_1, B_2, \dots, B_n) =$$

$$= \begin{bmatrix} B_1 & & & & \\ & B_2 & & & \\ & & B_3 & & \\ & & \dots & & \\ & & & B_n & \end{bmatrix}$$

$n \in N \cup \{\infty\}$.

Comment 8 Note the notation identification: $\zeta_F = \exp_{\odot}[\mathbf{A}_F] \equiv (1 - \mathbf{A}_F)^{-1 \odot}$. Note that $n! = 1 \text{ mod } 2$. Colligate also aside (?) reason:

$$\lim_{q \rightarrow 1} \exp_q = \exp$$

while

$$\lim_{q \rightarrow 0} \exp_q[x] = (1 - x)^{-1}.$$

Consult the Remark in [1] on the cases: Boolean poset 2^N and the "Ferrand-Zeckendorf" poset of finite subsets of N without two consecutive elements.

Observation 5 *The F -poset $P(G) = (\Phi, \leq)$ or equivalent to say: its cover relation graded digraph $G = (\Phi, \prec \cdot) = \bigoplus \rightarrow_{k=0}^m G_k$ **textcolorblue****is of Ferrers dimension one iff** in the process of deleting arcs from the cobweb poset Hasse diagram $D = (\Phi, \prec \cdot) = \bigoplus \rightarrow_{k=0}^n \langle \Phi_k \rightarrow \Phi_{k+1} \rangle$ does not produces 2×2 permutation submatrices in any of bigraphs G_k biadjacency matrices $B_k = B(G_k)$, $k = 0, \dots, n$, $n \in N \cup \{\infty\}$.*

3 Cobweb posets and differential posets of Stanley [3,4]

3.1. Preliminaries

The graded digraph G (graph) is utterly denominated by its sequence of bipartite digraphs (graphs) that every two consecutive levels of G do constitute.

The complete graded digraph D is utterly denominated by its sequence of complete bipartite digraphs - di-bicliques [1] that every two consecutive levels of D do constitute (see KoDAGs in [1] and references therein, consult the Example 2.7.2 in [5]).

Because of their appearance, an at a first glance outlook - these complete graded digraph D associated posets were called cobweb posets [digraphs D are identified with Hasse diagram of cobweb posets see KoDAGs in [1,2] and references therein).

Comment 9.

The appearance of "almost complete" graded digraph (subgraphs of KoDAGs) is tremendously prevailing. These look like - the hoary tree with silver cobweb threads. The other extreme to the complete in such a picture of a tree with cobweb (KoDAG) is an also beautiful melancholic bare rooted directed tree graph - void of this spider's web hoary tunicate and droplets (loops).

Here are come some examples listed with the convention that graphs become digraphs with all arcs directed upward (in the direction of increasing rank) or dually - downwards (in the direction of decreasing rank). If so has been done these become graded DAGs. First let us establish-recall for clarity that the directed tree (all arcs directed away from its root) is a digraph which becomes a tree if directions on the edges are ignored. Colligate with an arborescence. Naturally every arborescence is a directed acyclic graph.

Now come examples of sub-cobweb posets digraphs (hence DAGs) [5,3,4].

0. The binary tree digraph (directed Tree)
1. The Fibonacci digraph ([1], directed Tree)
2. The Young graph
3. The Young-Fibonacci graph
4. The Young-Fibonacci insertion graph
5. The 2-dimensional Pascal graph

6. The lattice of binary trees graph
7. The lattice of Bracket tree graph
8. The Fan_k graph [5] (Amalgamate k disjoint infinite chains by gluing their roots ("zeros"))
9. The special complete graded graph [5] (the set of vertices the same as in 8., connect any two vertices from consecutive levels Φ_k and Φ_{k+1} with an upward directed arc)
10. The F - denominated (hence any) complete graded digraph with finite minimal elements set (KoDAG in [1,2] and references therein).

3.2. Cobweb posets and differential posets

The class of posets known as differential posets were first introduced and studied by Stanley in [3] then [4]. A differential poset from [3] is a partially ordered set with raising and lowering operators U and D which satisfy the commutation relation $DU - UD = rI$ for some integer $r > 0$. Generalizations of this class of posets were studied by Stanley [4] and Fomin [5]. A number of examples of generalized differential posets are given in these [3,4,5] papers. Another example, a poset of rooted unlabelled trees, was introduced by Hoffman [6].

Let us consider at first the case $r = 1$ which we shall call GHW case for the reasons to become apparent soon.

In this GHW $r = 1$ case the operators U and D are defined correspondingly, $(x, y, z \in \Phi)$:

Definition 7

$$Dx = \sum_{y \prec \cdot x} y$$

$$Ux = \sum_{x \prec \cdot y} y$$

extended by linearity to (say complex, or...) linear space $C[\Phi]$.

From Theorem 2.2. in [3] we then have (consult also [7]) that GHW commutation relation $DU - UD = I$ holds iff $P(D) = \langle \Phi, \leq \rangle$ is differential ($r = 1$) poset. Out of this one infers inductively [8,9,10,11,7] what follows.

Observation 6

$$DU^n = nU^{n-1} + U^n D,$$

for $n \in N$.

Observation 7 *Cobweb posets Hasse digraphs from 10. above are examples of $\vec{q}$, $\vec{r}$ - differential posets (see [5] for all $q_n = 1$) and might serve for more general structures (due to Fomin [5]) called Dual graded Graphs.*

Check. Indeed. In [5] U_n and D_n are defined as restrictions of U and D onto $C[\Phi_k]$ "homogeneous" subspaces of $C[\Phi]$, $k = 0, 1, 2, \dots$. Then

$$D_{n+1} = q_n U_{n-1} D_n + r_n I_n, \quad n \in N,$$

where $r_0 = 1_F, q_0 = 0$ for $n = 0$ and $q_n = \frac{(n+1)_F}{(n-1)_F}, r_n = 0$ for $n > 0$.

For Fomin examples 8. and 9. above (see 2.7.2. in [5]) the (1.4.11) i.e. the (1.4.10) with all $q_n = 1$ from [5] holds for $r_0 = t \equiv 0_F, r_1 = r_2 = r_3 \dots = 0$

The KoDAGs graded digraphs example 10. in view of Observation 7 and Observation 6 becomes the motivating example of the following description appealing to the corresponding R. Stanley definition from [3]. One expect this description to be efficient and tangible while the combinatorics of path counting is concerned also via colligation with quantum models - for example.

Definition 8 Let F be such that $0_F = 1$. The locally finite graded poset $\Pi = \langle \Phi, \leq \rangle$ is then said to be F -differential poset iff

1. if $x, y \in \Phi, x \neq y$ and there are k_F elements of Φ which cover both of them then there are exactly k_F elements of Φ which are covered by both of them.
2. If $x \in \Phi$ covers exactly k_F elements from Φ then this very x is covered exactly by $(k + 1)_F$ elements from Φ

As in $F = N$ case (i.e for all $k = 1, 2, \dots$ we have $k_F \equiv k_N \equiv k$) the above conditions determine the number of elements and cover relations among them in the up to fourth rank [7]. Above the fourth rank ($k > 4$) cover relations and number of elements in levels Φ_k vary. And what does happen when the requirement $0_F = 1$ is relaxed?

Let us introduce the direct sum of certain projection on Φ_n operators and denote this operator with the symbol δ_F . Let x_n denotes homogeneous element of Φ i.e $x \in \phi_n$. Then δ_F is specified as follows.

Definition 9 $\delta_F = \text{diag}(1_F - 0_F, 2_F - 1_F, 3_F - 2_F, \dots, (n + 1_F - n_F, \dots) \equiv \text{diag}(\delta_0, \delta_1, \delta_2, \dots, \delta_n, \dots)$ i.e. $\delta_F(x_n) = \delta_n x_n$.

A straightforward verifying (see [7]) leads us to thus confirmed conclusion below; (note that $\delta_N = I$).

Observation 8

$$DU - UD = \delta_F$$

Out of this (as in [10,11]) one infers inductively what follows.

Observation 9

$$DU^n = n\delta_F U^{n-1} + U^n D,$$

for $n \in N$.

Since $Dx_0 = 0$ we have $DU = n\delta_F U^{n-1}x_0$ which means that D is representative of the Markowsky general linear operator i.e. a derivative from extended umbral calculus (see [10,11,12] and plenty of references therein).

The F -denominated cobweb posets are in a sense a canonical example of F -differential posets. More on that is expected soon.

Comment 10 (miscellaneous (aside?) final remark)

The ingenious ideas of differential and dual graded posets that we owe to Stanley

[3,4] and Fomin [5] bring together combinatorics, representation theory, topology, geometry and many more specific branches of mathematics and mathematical physics thanks to intrinsic ingredient of these mathematical descriptions which is the Graves - Heisenberg - Weyl (GHW) algebra usually attributed to Heisenberg by physicists and to Herman Weyl by mathematicians and sometimes to both of them (see: [3] for Weyl, [5] for Heisenberg and then [8] and [9] ; for GHW see [10-12] then note the content and context of [13,14]). As noticed by the author of [9] the formula

$$[f(a), b] = cf'(a)$$

where

$$[a, b] = c, [a, c] = [b, c] = 0$$

pertains to Charles Graves from Dublin [8]. Then it was re-discovered by Paul Dirac and others in the next century.

Let us then note that the picture that emerges in [11-12] discloses the fact that any umbral representation of finite (extended) operator calculus or equivalently - any umbral representation of GHW algebra makes up an example of the algebraization of the analysis with generalized differential operators acting on the algebra of polynomials or other algebras as for example formal series algebras.

Bibliography remark. On Umbra Difference Calculus references streams see [15] including references ad Comment 8 and all of that. On the history of cobweb poset ζ function formulas and also for the ζ_{-1} formula see [16].

References

- [1] A.K. Kwaśniewski , *Cobweb Posets and KoDAG Digraphs are Representing Natural Join of Relations, their di-Bigraphs and the Corresponding Adjacency Matrices*, arXiv:math/0812.4066v1,[v1] Sun, 21 Dec 2008 23:04:48 GMT
- [2] A.K. Kwaśniewski , *Some Cobweb Posets Digraphs' Elementary Properties and Questions* arXiv:0812.4319v1, [v1] Tue, 23 Dec 2008 00:40:41 GMT
- [3] Richard P. Stanley, *Differential posets*, J. Amer. Math. Soc. 1 (1988), 919-961.
- [4] Richard P. Stanley, *Variations on differential posets, Invariant Theory and Tableaux* (D. Stanton, ed.), Springer-Verlag, 1990, pp. 145-165.
- [5] Sergey V. Fomin *Duality of graded Graphs*, Journal of Algebraic Combinatorics, Volume 03, Number 4, October 1994 , pp. 357-404.
- [6] M. E. Hoffman, *Combinatorics of rooted trees and Hopf algebras*, Trans. Amer. Math. Soc. 355 (2003), 3795-3811.
- [7] Lewis, Joel B. *On Differential Posets, (2007) On Differential Posets*. Undergraduate thesis, Harvard College. <http://www.hcs.harvard.edu/thesis/repo/13/>

- [8] Charles Graves, *On the principles which regulate the interchange of symbols in certain symbolic equations* Proc. Royal Irish Academy vol. 6 , 1853-1857 , pp. 144-152
- [9] O.V. VISKOV, *On one result of George Boole*, (in Russian) Integral Transforms and Special Functions Bulletin 1 (1997) 2-7.
- [10] A.K.Kwaniewski *Extended finite operator calculus - an example of algebraization of analysis* Bulletin of the Allahabad Mathematical Society Vol 20 2005 pp. 1-24 arXiv:0812.5027v1, [v1] Tue, 30 Dec 2008 08:09:29 GMT
- [11] A.K.Kwaniewski *Main theorems of extended finite operator calculus*, Integral Transforms and Special Functions Vol. 14, No 6, pp.499-516, 2003
- [12] A.K.Kwaniewski *More on Bernoulli-Taylor formula for extended umbral calculus*, Advances in Applied Clifford Algebras, Vol. 16 (1) (2006) 29-39. arXiv:0802.1690v1, [v1] Tue, 12 Feb 2008 19:24:00 GMT
- [13] A. Dimakis, F Mller-Hoissen and T Striker *Umbral calculus, discretization, and quantum mechanics on a lattice* J. Phys. A: Math. Gen. **29**, (1996) 6861-6876
- [14] P .Blasiak, G. H. E. Duchamp, A. Horzela, K. A .Penson and A. I .Solomon, *Heisenberg-Weyl algebra revisited: combinatorics of words and paths* 2008 J. Phys. A: Math. Theor. **41**, 415204
- [15] A.K.Kwaniewski *First Contact Remarks on Umbra Difference Calculus References Streams* Bull. Soc. Sci. Lett. Lodz, **60**, (2005) 17-25. ArXiv: math.CO/0403139 , [v1] Mon, 8 Mar 2004 21:12:56 GMT
- [16] A.K. Kwaśniewski , *Graded posets inverse zeta matrix formula* arXiv:0903.2575v1 [v1] Sat, **14 Mar 2009** 20:32:01 **130-th Birthday of Albert Einstein**